

**A New Model for Understanding the Role of Environmental Factors in the Origins of
Chronic Illness:
A Case Study of Type 1 Diabetes Mellitus**

Medical Hypotheses, Vol 63, Issue 6, 2004. pp. 1035-1046

Elsevier B. V., Science Direct
<http://www.sciencedirect.com/mehy>

Veronique P. Mead, MD, MA

ABSTRACT

There is a need for a new pathophysiological model explaining and linking the role of non-genetic factors believed to contribute to origins of many chronic physical diseases. This article presents a theoretical model for explaining the confusing and often contradictory findings regarding the role of environmental influences in chronic illness. A case study of type 1 diabetes is presented to illustrate the applicability of this model. Type 1 diabetes is a disease that has been widely studied, for which clear diagnostic criteria exist, and for which development of effective prevention strategies represents significant challenges.

The model is formulated from the large database of research on our increasing understanding of the interaction between environmental factors, physiology, and autonomic regulatory function. The model integrates data from research in the fields of the experience-dependent maturation of the nervous system and the neurophysiology of traumatic stress. This article describes how disruptions in early bonding and attachment, including adverse events such as traumatic stress, are capable of causing 1) long-term imbalances in autonomic regulatory function and 2) relative dominance of sympathetic or parasympathetic activity.

The proposed model of autonomic dysfunction suggests that ongoing mechanisms promoting high glucose in the context of decreasing insulin production in type 1 diabetes represents a state of relative sympathetic dominance. It is further proposed that this state is influenced by environmental factors, which affect autonomic, immune and endocrine systems during critical period programming. The model also identifies a link between the many seemingly unrelated non-genetic risk factors, and appears capable of explaining contradictions and enigmas in epidemiological and clinical studies regarding non-genetic origins of type 1 diabetes, including the role of stress, variation in age of onset, and duration of the preclinical phase. The case study and model are also helpful for understanding similar environmental contributions to origins of other chronic illnesses, such as multiple sclerosis, rheumatoid arthritis and Parkinson's disease.

Table of Contents

INTRODUCTION 3

I. Non-genetic factors affecting ANS function..... 4

1. The experience-dependent maturation of the nervous system..... 4

 Gene-environment interactions during development 4

 Synapse development 4

 Critical period programming..... 4

 Influence of the prenatal environment..... 4

 Maternal-infant bonding & psychobiological regulation 4

 The sensitive period..... 5

 Bonding disruption 5

2. Traumatic Stress..... 6

 Definition and physiology 6

 Risk for ANS dysfunction 6

 Kindling 6

 Reinstatement..... 6

 The role of perception 7

 Risk in early life 7

 Prenatal stress..... 7

 Rate of progression..... 8

III. A Model of Autonomic Dysfunction..... 8

IV. Relevance of the model to non-genetic risk for type 1 diabetes..... 8

 Stress, traumatic stress & type 1 diabetes 8

 Stress and glucose variations in diabetes..... 9

 Risk factors preceding diagnosis..... 9

 Perinatal risk factors..... 10

 Early separation and stress represent common denominators linking perinatal risk factors 10

 Breastfeeding and cow’s milk..... 11

 Timing of diagnosis..... 11

 Age of onset 11

 The preclinical phase..... 11

 Rate of progression..... 11

 Autoimmune activity..... 11

CONCLUDING REMARKS 12

Acknowledgements 12

REFERENCES 13

A New Model for Understanding the Role of Environmental Factors in the Origins of Chronic Illness

INTRODUCTION

One of the unexplained pathophysiological processes associated with type 1 diabetes is the seemingly contradictory secretion and production of glucose from multiple physiological mechanisms, such as gluconeogenesis, glycogenolysis and increased secretion of glucagon, which occur in the face of existing elevations in glucose. Despite the fact that relative or absolute insulin deficiency is considered to be the cause of increases in glucose associated with type 1 diabetes, lack of insulin does not account for this aspect of hyperglycemia.

Although peptides, neurotransmitters, and other nutrients affect insulin secretion, glucose is the key regulator¹. Glucose is the primary and preferred source of energy for the body and is the major and ultimate source for nerve cells², which do not require insulin for the utilization of glucose³. Because glucose is so vital to the brain, and because the brain cannot store glucose, the nervous system is intricately geared to maintain a minimum critical level of glucose in the blood. While many hormones influence glucose levels and storage, glucose metabolism is ultimately regulated by the nervous system^{4,5}.

The nervous system regulates not only glucose activity but also interacts with and regulates the immune system. Although type 1 diabetes is described as resulting from an autoimmune process, it is also categorized as a disease of autonomic dysfunction because of the nature of many of the long-term complications¹.

Non-genetic factors such as stress influence nervous system activity,

including the capacity to modulate the regulation of glucose metabolism⁴. Environmental factors are believed to account for greater than 50% of the risk for type 1 diabetes^{6,7} and have been hypothesized to unmask the disease in individuals with a genetic predisposition^{1,8-10}. This article builds on research describing the influence of environmental factors on the nervous system and presents a new model for explaining the influence of non-genetic factors on the origins of type 1 diabetes. The model integrates neurophysiological findings from the developmental and psychophysiological literature, drawing from two relevant fields of study: 1) the experience-dependent maturation of the nervous system, in which the environment of early experience influences the structural and functional development of the nervous system, and 2) traumatic stress, which is increasingly understood to be capable of exerting complex and frequently long-lasting effects on activity of the autonomic nervous system (ANS). These fields of study have been reviewed elsewhere¹¹⁻¹⁵ and important concepts are summarized below to provide a foundation for understanding current perspectives on factors known to influence states of autonomic dysfunction. These principles are then applied to provide a common denominator explaining a link between the varied and seemingly unrelated environmental risk factors for type 1 diabetes¹⁶.

I. Non-genetic factors affecting ANS function

1. The experience-dependent maturation of the nervous system

Gene-environment interactions during development

The nervous system is immature at birth¹⁷ and maturation takes place in early life during relatively defined periods of time. Pruning of synapses is genetically timed¹⁸ and the number^{19,20} as well as strength^{13,21} of synapses is influenced by interactions with the environment^{11-13,22}. The contribution of environmental factors is referred to as experience-dependent maturation^{12,13,20}.

The process of experience-dependent maturation: 1) influences the density of synapses in different tissues as well as variations in density in the same tissues in different individuals²³, 2) can predispose to relative predominance of sympathetic or parasympathetic activity, and 3) fosters unique individual responses to different types of stress¹¹. This process promotes plasticity of the nervous system and facilitates human organisms' finely honed capacity for adaptation to their own unique environments^{12,13}.

Synapse development

Cortical synaptogenesis begins prenatally, peaks by approximately 1 to 2 years of age, and appears to be activity-dependent²⁴. Genetically timed apoptosis and elimination of synapses follows and is at least partly environmentally regulated²⁴. Timing of cortical synapse elimination varies in different areas, is most dramatic from one year of age through mid-adolescence^{24,25}, and continues at a slower

rate in adulthood^{20,24}. Imaging studies used to evaluate growth rates of the developing nervous system have tracked patterns of glucose utilization as a measure of metabolism, synaptogenesis, and plasticity to demonstrate that growth is particularly active between 4 and 9 to 10 years of age, when glucose utilization is at its highest²⁶. Levels of growth then decline to reach adult values by the ages of 16 to 18²⁶.

During development, nerve pathways that are reinforced, such as through frequent utilization, are generally promoted through the stabilization and strengthening of synapses, while those that are underutilized are reduced through selective pruning¹². Gene-environment interactions contribute to individuality, take place both in prenatal and postnatal life^{12,27,28}, and are believed to influence risk for pathology and disease^{11,23,29,30}.

Critical period programming

The timing of environmental events most strongly influences developing structures^{11,28} and has the highest impact on organ systems undergoing periods of rapid growth^{11,31}. Critical period programming occurs primarily in prenatal and early life. Adverse events occurring during these times may have long-term and irreversible effects on the developing organism^{20,23}.

Influence of the prenatal environment

Maternal-infant bonding & psychobiological regulation

The experience-dependent maturation of the nervous system is affected by interactions with the environment in general, and by the attachment bond between infant and primary caregiver(s) in particular^{11-13,32,33}.

The mother, who's role has been the most frequently studied in the function of primary caregiver^{11,12}, serves as a "psychobiological regulator"¹¹ for her dependent and essentially helpless infant¹². In this capacity, she helps to modulate his or her levels of arousal to facilitate the establishment of self-regulation not only of behavioral rhythms, but also of physiological rhythms, including autonomic, neurochemical, and hormonal functions¹¹.

Many of the interactions that influence the ANS and the balance between parasympathetic and sympathetic activity occur at an unconscious nonverbal level³⁴ through a multitude of interactions inherent to parent-infant interactions, including holding, gazing, and soothing¹¹. The mother's ability to respond to and to stimulate her infant at optimal levels is influenced by the degree of attunement with her infant, and serves to buffer his or her physiological^{12,35} as well as emotional and behavioral responses to stress¹¹.

Attunement between mother and child is directly affected by the maternal-infant bond, which in turn is shaped by prenatal and perinatal events^{11,36}. Among the complex factors that influence bonding at birth are the mother's attitude toward the pregnancy and her perception of available support systems^{36,37}, her experience of procedures such as amniocentesis^{38,39}, and her perception of stress during pregnancy^{36,40,41}.

The sensitive period

Among the most influential perinatal experiences affecting bonding are maternal-infant interactions in the hours and weeks following birth. Early contact during the first day in general, and the first hour postpartum in particular, appears to be of special importance^{36,42}. During the

first hour the newborn exhibits qualities of alertness and exploratory behavior that do not occur again to the same extent for several weeks³⁶. Contact during this first hour has been found to increase the number of mothers that breastfeed, the duration of breastfeeding⁴³, or both^{36,44,46}. In addition, early contact also appears to improve the quality of future behavioral interactions between mother and infant^{42,44,47-49} and to reduce the frequency of early infections in the baby during the first months of life³⁶. Measurable effects have been noted in the quality of maternal-infant interactions and infant development up to one⁵⁰ and three years⁴⁶ of age following early contact.

Bonding disruption

Separation in early life is associated with changes in hypothalamic-pituitary-adrenal (HPA) responses to stress⁵¹, transient and long-term changes in immune competence in non-human primates⁵², and reduced maternal-infant attunement⁵³. The impact of maternal-infant separation during the sensitive period may permanently alter affectional ties³⁶, and may consequently influence developing organ systems, including the nervous system¹¹. Events that affect the ability of the mother to attend to her infant shape the capacity of the newborn to tolerate stress, since the immature nervous system is unable to regulate states of high arousal. Events occurring during labor and delivery that may affect the mother or the infant's ability to bond include early separation, pain in the mother or infant, the use of medication such as anesthesia, and anxiety, among others³⁶.

Whereas healthy newborns demonstrate more rapid returns to baseline cortisol following exposure to stress⁵⁴, babies born following mild obstetrical

complications have less optimal HPA responses⁵⁵ as well as decreased habituation and sensitization to stressors⁵⁴. Maternal-infant separation following cesarean sections is common and appears to negatively impact quality of maternal-infant interactions⁵⁶⁻⁵⁹ as well as frequency of breastfeeding⁵⁹.

2. Traumatic Stress

Definition and physiology

The definition of traumatic stress has been summarized as the *experience of an event that is perceived by a relatively helpless individual to be both life threatening and inescapable*^{15,60}. Although the delineation between different types of stress has been difficult to fully elucidate, useful factors in the characterization of traumatic stress are that it is associated with extreme fear and high emotional charge^{15,61,62}.

Highly emotional experiences promote the imprinting of an event⁶³, along with associated contextual environmental and physiologic cues, into procedural (unconscious) memory^{15,64}. In comparison with other types of conditioning, fear conditioning can occur following one trial and can be extremely long-lasting¹⁵. The imprinting of highly charged experiences creates conditioned responses that even when extinguished remain stored in memory and available for activation following exposure to an environmental trigger⁶³. These characteristics of the nervous system represent innate survival strategies that serve as internal references to promote survival by enabling the organism to respond to future experiences with enhanced rapidity and efficiency¹¹.

Traumatic stress is capable of reprogramming the HPA axis and affects

ANS regulatory function, resulting in states of exaggerated autonomic cycling and arousal⁶⁵, especially in response to stress¹⁵. Traumatic stress can also result in states of prolonged sympathetic arousal^{15,66} as well as disruption in the interdependence between the two branches of the ANS¹¹.

Risk for ANS dysfunction

Kindling

The term kindling dates from studies of rats in which the repeated use of low intensity electrical stimulus produces or kindles seizures that become self-perpetuating and that require no further stimulus for seizure activity to occur^{64,67}. The seizures are the result of new circuits that are activated and kindled¹⁵ by physiologic and psychological responses to internal and external cues^{64,68} increasingly distinct from the original learned trigger or conditioned stimulus⁶³. The conditioned responses that imprint during traumatic stress appear to be mediated by changes in the hippocampus occurring at the time of high arousal¹⁵. They may also take place through the creation of new cell assemblies, in which neurons that initially fire spontaneously together in response to certain stimuli are more likely to eventually fire without obvious stimulation, a Hebbian model referred to by Ledoux⁶³. The concept of *kindling* is utilized in the neurological literature^{15,64,67} and in models of traumatic stress^{69,70}.

Reinstatement

Conditioned responses occurring as a consequence of traumatic stress may be quiescent or inaccessible for long periods of time, but appear to remain in procedural memory⁶³. Following exposure to an

event similar to the original stimulus⁷¹ or to some unrelated stressful experience^{63,64} these memories can be kindled and the conditioned responses “reinstated”. Conditioned responses associated with PTSD such as hyperarousal become activated with increasing ease following exposure to adverse environmental cues. The effects of traumatic stress are consequently considered to be intrinsically self-perpetuating^{15,70}.

The role of perception

Although certain types of events such as earthquakes or the witnessed murder of a loved one are more likely to be experienced as traumatic because of their overwhelming nature, it is the perception of an event rather than the specific nature of an event that influences kindling, reinstatement, and risk for ANS dysregulation^{15,60}. Risk is influenced by a number of factors, including intensity⁷² and degree of life threat perceived during the original traumatic stressor⁷³, previous experience of trauma⁷⁴, and subsequent exposure to seemingly irrelevant as well as related stressors⁶¹. ANS function is also affected by the perceived availability and quality of social supports^{61,75} and buffers⁶¹, history of bonding and attachment^{15,61,73}, and stage of emotional and physiological development, all of which influence the capacity to regulate arousal and to cope with stress⁷⁶. Because experiences are unique to each individual, the determination of whether an event is experienced as traumatic, as well as risk for kindling, reinstatement, symptomatology, and disease^{11,77} appears to be idiosyncratic¹⁵.

Risk in early life

Stress in early life influences the immune and nervous systems, among

others⁷⁸. Children are at increased risk of perceiving stressful events as intense and life threatening because of their inherent dependence and relative helplessness¹⁴, and are consequently particularly vulnerable to PTSD¹⁵. Risk for adverse physiologic consequences is believed to be due to the high degree of nervous system development²⁰ that occurs during childhood and to the fact that immature nervous systems are incapable of regulating states of high arousal¹¹. A strong bond has been found to promote while a disrupted bond to interfere with the ability of a child to regulate arousal⁷⁹ and early separation from parents has been identified as an important risk factor for PTSD⁸⁰. The experience of traumatic stress in the context of relationship with another human being is one of the most influential risk factors for ANS dysregulation and PTSD^{15,61}. Secure attachment, on the other hand, is one of the most protective factors against the development of symptoms following exposure to traumatic stress¹⁴.

Prenatal stress

Early exposure to non-genetic factors such as stress in prenatal life stimulates the fetal HPA axis⁸¹, can permanently affect the number and sensitivity of glucocorticoid receptors, and can program the HPA axis for life^{31,81}. Number of glucocorticoid receptors has been found to be proportional to the severity of symptoms of PTSD⁸². Maternal exposure to prenatal stress has also been found to predict birth size and gestational age independent of biomedical risk⁸³⁻⁸⁵, and to influence physiological as well as psychological development postpartum³⁷. Size at birth appears to be influenced by the timing⁸³ and quality^{81,83} of emotional stress experienced by the mother during

pregnancy, as well as by her perceived availability of social support⁸⁴.

Rate of progression

The rate at which reinstatement occurs following traumatic stress is idiosyncratic and symptoms of PTSD can occur weeks, months or years following the original event^{15,61}. High intensity initial and subsequent exposure to relevant stressors promotes a more rapid response of neurophysiological hyperreactivity¹⁴ and increases the rate with which reinstatement can occur. Exposure to insufficient stimuli may allow symptoms to resolve after reinstatement or enable a dormant conditioned response to remain inaccessible⁶³. Although the development, progression, and resolution of symptoms of PTSD are variable and occur in relatively few individuals following traumatic stress⁸⁶, symptoms are often remarkably refractory to treatment once they develop⁶¹.

III. A Model of Autonomic Dysfunction

The model proposes that the pathophysiology of type 1 diabetes is related to altered autonomic regulation of glucose activity. While complex, the specific nature of ANS dysfunction is suggested to be influenced by a predominance of sympathetic activity or tone, a physiological state that promotes glucose availability in the face of simultaneous inhibition of insulin secretion and activity. Dysregulation of the sympathetic branch of the ANS has long been hypothesized to play a role in the pathogenesis⁸⁷ and symptomatology^{88,89} of type 1 diabetes, although the causes of such dysregulation have not been understood.

The model proposes that dysfunction of the sympathetic nervous system associated with type 1 diabetes is at least partly influenced by exposure to adverse events, including traumatic stress, during the experience-dependent maturation of the nervous system. Initial exposure to such events is suggested to occur during critical period programming of structures relevant to glucose and insulin regulation, such as the nervous system. Kindling and reinstatement of SNS patterns of dysfunction are proposed to occur following sufficient exposure to relevant stressors during periods of programmed synapse pruning and growth, which predominates during preschool age and adolescence. At-risk individuals may fail to develop diabetes or may do so at later ages if they are not exposed to relevant types, number, or intensities of stressors. This is hypothesized to occur because the predisposing conditioned responses of autonomic SNS dysfunction fail to kindle or kindle more slowly.

IV. Relevance of the model to non-genetic risk for type 1 diabetes

Stress, traumatic stress & type 1 diabetes

Stress has long been suggested^{1,90-94}, and recently confirmed as a risk factor for type 1 diabetes⁹⁵. Many reports exist describing the onset of diabetes within days or weeks of intense emotional events⁹⁶⁻⁹⁸ and traumatic stress was believed by many to be an important risk factor for type 1 diabetes in the first half of the twentieth century⁹⁴. One means by which traumatic stress was believed to cause diabetes was by reactivation of infantile neuroses that were indirectly expressed through the vegetative (autonomic)

nervous system^{92,98,99}. Models of traumatic stress as a cause of diabetes fell into disfavor, however, when a smaller number of soldiers than anticipated developed the disease following service in World War I and II⁹⁴ and direct causal links with psychogenic factors could not be made¹⁰⁰.

The proposed model provides a framework for explaining the role of stress and trauma in the etiology of type 1 diabetes and suggests that individuals who develop the disease are those who have been exposed to traumatic stress during a specific time in early life *and* who have experienced appropriate intensity and frequency of subsequent stressors sufficient to promote kindling and reinstatement. From this perspective, the small number of soldiers who developed diabetes would have been the few who had experienced both relevant traumatic experiences during critical period programming in early life so as to be predisposed to patterns of SNS dysfunction⁹² and who were also subsequently exposed to sufficient and appropriately timed relevant idiosyncratic stressors.

Stress occurring in the year before diagnosis has been found to be the only variable affecting age of onset in humans in one large study and has been proposed to precipitate rather than initiate the disease¹⁰¹. Stress has also been found to reduce the age of onset of diabetes in BB rats¹⁰².

Stress and glucose variations in diabetes

The frequent finding of glycemic variation (or lack thereof) in response to stress among individuals with type 1 diabetes¹⁰³⁻¹⁰⁵ appears to be idiosyncratic¹⁰⁶⁻¹⁰⁸. These variations are consistent with models of kindling for conditioned responses to idiosyncratic physiological, psychological or other environmental cues.

Risk factors preceding diagnosis

A variety of factors occurring in the years preceding diagnosis of type 1 diabetes have been proposed to influence risk. In addition to stress^{90,93,109,110} and trauma^{94,96}, risk factors include infection^{9,95}, a cold environment⁸, and seasonality¹¹¹⁻¹¹⁴, to name a few. Studies attempting to clarify the relationship between identified factors and type 1 diabetes have frequently demonstrated conflicting results, however. For example, infection is associated with increased risk for type 1 diabetes in some studies^{115,116}, but has been found to be unassociated or even protective in others^{117,118}.

According to the model, it is not the specific nature of an event that represents risk for diabetes, but the idiosyncratic nature in which events are perceived. Such a perspective is consistent with the proposal that stress is the common link between perinatal risk factors for type 1 diabetes as well as risk factors in the period preceding diagnosis¹¹⁹. This model predicts contradictions in the identification of risk factors.

Seasonal risk for type 1 diabetes, which is increased in fall and winter in certain age groups⁹, may represent a stressful event and result from increased workload on beta cells¹⁰⁹. Physiologic increases in SNS activity occur in mammals prior to and during cold seasons as a result of changing energy needs, glucose utilization and storage, and immune activity¹²⁰. Changes in sympathetic activity associated with seasonal variation may promote kindling and reinstatement in individuals predisposed to states of increased sympathetic tone.

Perinatal risk factors***Early separation and stress represent common denominators linking perinatal risk factors***

Important environmental risk factors for type 1 diabetes appear to originate during a limited period of exposure¹²¹ in early life^{1,109,122-124} and are proposed to be factors to which only children are exposed¹²¹. These environmental risk factors probably occur during prenatal life^{7,90,109,121,125-127} and include maternal-related events¹²¹. Perinatal risk factors proposed for type 1 diabetes include maternal infection^{9,126,128}, respiratory disease in the newborn^{119,123}, preeclampsia^{119,123,127}, cesarean section^{119,124,127}, maternal-child blood group incompatibility and jaundice^{119,123}, amniocentesis and labor complications¹²⁷, shorter gestational age and larger birth size¹¹⁹, and certain patterns of intrauterine growth¹²⁹, among others. According to the proposed model, early events affect risk for type 1 diabetes through complex interactions between protective and adverse events that influence experience-dependent maturation of the nervous system. A common denominator among perinatal risk factors for type 1 diabetes is an increased risk for disruptions in maternal-infant bonding, such as occur following stress and early separation.

Intrauterine growth patterns, larger size at birth, and shorter gestation have been proposed to reflect prenatal exposure to non-genetic events¹²⁹ such as stress³¹. Prenatal stress in the mother and prenatally, for example, may be experienced during many of the identified perinatal risk factors for type 1 diabetes, including procedures such as amniocentesis^{41,130}. The experience of stress may precede the procedure, vary according to the reasons for the test, and

last until results have been received. Positive results may be reassuring to the mother and may facilitate bonding whereas abnormal results may impact stress levels as well as bonding³⁸.

Early separation was found to be the only factor identified with risk for type 1 diabetes in an animal study evaluating risk associated with jaundice with or without phototherapy¹³¹. Other identified perinatal risk factors also appear to increase risk for early maternal-infant separation, particularly during the sensitive period immediately following birth. As previously discussed, even brief early separation influences bonding, capacity for physiological self-regulation in the infant, and experience-dependent maturation, and may be perceived as traumatic by the newborn.

Exposure to maternal states of sympathetic arousal during experience-dependent maturation in the first three years of life contributes to predisposition for states of increased sympathetic tone in the infant and in the growing child¹¹. Such effects may last into adulthood¹¹. Risk factors such as preeclampsia, which has been associated with states of increased sympathetic activity in the mother¹³²⁻¹³⁶, may have similar effects, as may cesarean sections, which can be associated with maternal postpartum traumatic stress¹³⁷.

Early life events involve complex idiosyncratic interactions and predict variability in perinatal risk factors rather than strict consistency. The apparent lack of risk associated with some measures of perinatal distress such as reduced apgar scores¹¹⁹ may represent contributions from protective factors such as bonding.

Breastfeeding and cow's milk

Breastfeeding may be protective from risk for type 1 diabetes¹³⁸ and shorter duration^{95,101,138,139}, as well as lack of breastfeeding⁹⁰ are associated with increased risk in most studies. Contradictory findings, however, exist regarding a role for cow's milk in risk for type 1 diabetes¹⁴⁰⁻¹⁴³.

Higher rates of separation in early life are associated with decreases in the frequency and duration of breastfeeding³⁶. Decreases in duration of breastfeeding can reduce physiological and social bonding³⁶ and may interfere with protective factors that reduce risk for type 1 diabetes. The idiosyncratic nature of experience and complex interactions between risk and protective factors is consistent with contradictory findings regarding the role cow's milk¹⁴⁰⁻¹⁴³ and breastfeeding^{101,118,144,145} in risk for type 1 diabetes. According to the model, decreases in breastfeeding and the early introduction of cow's milk may represent disruptions in maternal-infant bonding¹⁴⁶.

Timing of diagnosis**Age of onset**

Incidence of type 1 diabetes is highest in children and adolescents¹⁴⁷, and peaks at 2, 3 to 6, and 10 to 14 years of age¹⁰. Only 25% of individuals develop type 1 diabetes after the age of 21¹⁴⁸.

The model suggests that life events associated with risk prior to the diagnosis of type 1 diabetes represent idiosyncratic stressors that kindle and reinstate existing conditioned responses of chronic sympathetic arousal. Risk factors are proposed to induce reinstatement during periods of genetically programmed synapse development, which predominate during experience-dependent maturation in early childhood and adolescence.

The preclinical phase

The preclinical phase leading to diabetes is frequently lengthy, lasting up to 10 years¹⁴⁹⁻¹⁵¹ or longer¹⁵². This period has been documented through the presence of autoantibodies as well as through changes in insulin levels, which have been shown to decline up to 11 years prior to the onset of diabetes¹⁵³.

The existence of a preclinical phase in diabetes has similarities with the latency period that frequently precedes onset of symptoms of PTSD. This pattern is consistent with the model, in which conditioned responses kindle during latency periods and are reinstated following exposure to sufficient idiosyncratic stressors in conjunction with synapse loss or proliferation.

Rate of progression

Although type 1 diabetes is commonly proposed to occur following exposure to certain environmental events preceding diagnosis, some investigators have proposed that predisposition occurs early in life and that variation in age of onset reflects different rates of disease progression rather than different ages of exposure to critical environmental events^{10,153}. The model supports the latter hypothesis.

Autoimmune activity

It has been proposed that non-genetic risk factors in early life initiate autoimmunity leading to type 1 diabetes while factors preceding diagnosis unmask or accelerate the disease process^{109,153}. The presence of autoantibodies to islet cells (ICA), insulin (IAA), and glutamate acid decarboxylase (GAD or GAA) is associated with increased risk for type 1 diabetes¹⁴⁸ even though none of these

antibodies are specific for beta cells^{1,154} and 10% of individuals have no detectable autoantibodies at diagnosis¹⁰.

Autoantibodies can be acquired early in life and develop over a period of months to years¹⁵⁵. They can fluctuate¹⁰, exhibit transiency^{156,157}, and have been found to remit in 10 to 78% of individuals, perhaps because the offending factor has been removed¹⁰. Only a small number of individuals with autoantibodies develop type 1 diabetes and highest risk is associated with early expression of IAA¹⁵⁸. Increased risk is also associated with persistence¹⁵⁶, sequential progression¹⁵⁵, and presence of multiple antibodies¹⁵⁸. The incidence of autoantibodies appears to decrease with age¹⁰ and conversion to autoantibody positivity is rare over the age of 10 years^{10,148,153}.

Although the mechanisms by which autoimmunity plays a role in type 1 diabetes remain unclear, the patterns of incidence, transience, and persistence of autoantibodies in relationship to risk for disease are consistent with models of experience-dependent maturation and traumatic stress. Activation and specificity of autoimmune processes may be regulated by ANS activity and promoted by states of dominant sympathetic activity, which persistently promotes inhibition of insulin and in some states also promotes increases in immune activity¹⁵⁹. Similarly, fluctuations in autoantibody levels may represent varying stages of kindling influenced by the frequency and quality of exposure to idiosyncratic stressors and buffers. It has been suggested that islet cell autoantibodies (ICAs) may represent secondary markers of islet cell destruction¹⁶⁰. Progression to disease in individuals

who are autoantibody positive has been proposed to depend on sufficient exposure to environmental pathogens or “multiple hits”¹⁶¹, a concept also consistent with kindling¹⁶².

CONCLUDING REMARKS

Our growing understanding of the interaction between environmental factors and human physiology provides powerful new perspectives from which to consider the origins of chronic physical disease such as type 1 diabetes. Such a model may be applicable to a developing a better understanding of the origins of other chronic diseases, ranging from type 2 diabetes, to multiple sclerosis and rheumatoid arthritis, to chronic fatigue, fibromyalgia, and multiple chemical sensitivity. The integration of concepts from multiple disciplines represents an exciting frontier that may contribute to identification of increasingly effective prevention and treatment strategies, and concepts presented in this model warrant further study.

Acknowledgements

Preparation of this manuscript was supported in part by a grant from the Diabetes Trust Foundation (formerly the Diabetes Trust Fund). The constructive comments of Robert Scaer, M.D. are gratefully acknowledged.

REFERENCES

1. E Braunwald, AS Fauci, DL Kasper, SL Hauser, DL Longo, JL Jameson, Eds: Harrison's principles of internal medicine. 15th ed. New York: McGraw-Hill; 2001.
2. WF Ganong: Review of medical physiology. Los Altos, CA: Lange Medical; 2001.
3. L Sherwood: Human physiology: from cells to systems. Minneapolis, MN: West Publishing; 1993.
4. K Nonogaki: New insights into sympathetic regulation of glucose and fat metabolism. *Diabetologia* **43** 5 (2000), pp. 533-549.
5. SC Woods, PJ Kulkosky: Classically conditioned changes of blood glucose level. *Psychosom Med* **38** 3 (1976), pp. 201-219.
6. Diabetes Epidemiology Research International: Preventing insulin dependent diabetes mellitus: the environmental challenge. *BMJ* **295** 6596 (1987), pp. 479-481.
7. D Kumar, NS Gemayel, D Deapen, D Kapadia, PH Yamashita, M Lee, JH Dwyer, P Roy-Burman, GA Bray, TM Mack: North-American twins with IDDM. Genetic, etiological, and clinical significance of disease concordance according to age, zygosity, and the interval after diagnosis in first twin. *Diabetes* **42** 9 (1993), pp. 1351-1363.
8. G Dahlquist: The aetiology of type 1 diabetes: an epidemiological perspective. *Acta Paediatr Suppl* **425** (1998), pp. 5-10.
9. KMM Alberti, P Zimmet, RA Defronzo, Eds: International textbook of diabetes mellitus. New York: John Wiley & Sons; 1997.
10. PH Bennet, MJ Rewers, WC Knowler: Epidemiology of diabetes mellitus. In: DJ Porte, RS Sherwin, editors. *Ellenberg and Rifkin's diabetes mellitus*. 5th ed. Stamford, CT: Appleton & Lange; 1997. pp. 373-400.
11. AN Schore: Affect regulation and the origin of the self: the neurobiology of emotional development. Hillsdale, NJ: Lawrence Erlbaum; 1994.
12. National Research Council and Institute of Medicine: From neurons to neighborhoods: the science of early childhood development. Committee on integrating the science of early childhood development. JP Shonkoff, DA Phillips, Eds. Board on children, youth, and families, Commission on behavioral and social sciences and education. Washington, D.C.: National Academy Press; 2000.

13. D Siegel: The developing mind. New York:NY: Guilford; 1999.
14. RS Pynoos, AM Steinberg, A Goenjian: Traumatic stress in childhood and adolescence: recent developments and current controversies. In: BA van der Kolk, AC McFarlane, L Weisaeth, editors. Traumatic stress: the effects of overwhelming experience on mind, body, and society. New York: Guilford; 1996. pp. 331-358.
15. RC Scaer: The body bears the burden: trauma, dissociation, and disease. New York: Haworth Medical; 2001.
16. VP Mead: Somatic psychology theory and the origins of chronic illness: a case study of type 1 diabetes [Master's Thesis]. Boulder (CO), 427 p: Master's Thesis, Naropa University; 2003 Available from author, 4439 Driftwood Place, Boulder, CO 80301 USA.
17. MA Hofer: Early stages in the organization of cardiovascular control. *Proc Soc Exp Biol Med* **175** 2 (1984), pp. 147-157.
18. JT Bruer: The brain and child development: time for some critical thinking. *Public Health Rep* **113** 5 (1998), pp. 388-397.
19. AM Turner, WT Greenough: Differential rearing effects on rat visual cortex synapses. I. Synaptic and neuronal density and synapses per neuron. *Brain Res* **329** (1985), pp. 195-203.
20. JE Black: How a child builds its brain: some lessons from animal studies of neural plasticity. *Prev Med* **27** (1998), pp. 168-171.
21. J Bruer: Neural connections: some you use, some you lose. Phi Delta Kappan. Volume accessed online 2003 April 11, http://www.jsmf.org/zarticles&pap/John/neural_connections.htm; 1999. pp. 264-277.
22. R Kotulak: Inside the brain: revolutionary discoveries of how the mind works. *Prev Med* **27** (1998), pp. 246-247.
23. JB Young, SF Morrison: Effects of fetal and neonatal environment on sympathetic nervous system development. *Diabetes Care* **21** (Suppl 2) (1998), pp. B156-160.
24. PR Huttenlocher, AS Dabholkar: Regional differences in synaptogenesis in human cerebral cortex. *J Comp Neurol* **387** 2 (1997), pp. 167-178.
25. PR Huttenlocher: Synaptic density in human frontal cortex-developmental changes and effects of aging. *Brain Res* **163** 2 (1979), pp. 195-205.

26. HT Chugani: A critical period of brain development: studies of cerebral glucose utilization with PET. *Prev Med* **27** (1998), pp. 184-188.
27. PR Huttenlocher: Synapse elimination and plasticity in developing human cerebral cortex. *Am J Ment Retard* **88** 5 (1984), pp. 488-496.
28. P Levitt, B Reinoso, L Jones: The critical impact of early cellular environment on neuronal development. *Prev Med* **27** (1998), pp. 180-183.
29. R Dubos, D Savage, R Schaedler: Biological Freudanism: lasting effects of early environmental influences. *Pediatrics* **38** 789-800 (1966), pp.
30. H Weiner: *Psychobiology and human disease*. New York: Elsevier; 1977.
31. P Nathanielsz: *Life in the womb*. Ithaca, NY: Promethean; 1999.
32. BO Gilbert, SB Johnson, J Silverstein, J Malone: Psychological and physiological responses to acute laboratory stressors in insulin-dependent diabetes mellitus adolescents and nondiabetic controls. *J Pediatr Psychol* **14** 4 (1989), pp. 577-591.
33. AN Schore: Attachment and the regulation of the right brain. *Attach Hum Dev* **2** 1 (2000), pp. 23-47.
34. MF Basch: The concept of affect: a re-examination. *J Am Psychoanal Assoc* **24** (1976), pp. 759-777.
35. MR Gunnar: Quality of early care and buffering of neuroendocrine stress reactions: potential effects on the developing human brain. *Prev Med* **27** 2 (1998), pp. 208-211.
36. MH Klaus, JH Kennell: *Maternal-infant bonding: the impact of early separation and loss on family development*. St. Louis: Mosby; 1976.
37. MJ Carey-Smith: Effects of prenatal influences on later life. *N Z Med J* **97** 747 (1984), pp. 15-17.
38. N Caccia, JM Johnson, GE Robinson, T Barna: Impact of prenatal testing on maternal-fetal bonding: chorionic villus sampling versus amniocentesis. *Am J Obstet Gynecol* **165** 4 Pt 1 (1991), pp. 1122-1125.
39. JW Spencer, DN Cox: A comparison of chorionic villi sampling and amniocentesis: acceptability of procedure and maternal attachment to pregnancy. *Obstet Gynecol* **72** 5 (1988), pp. 714-718.
40. TR Verny: Prenatal attachment. *J Obstet Gynecol Neonatal Nurs* **25** 6 (1996), pp. 464.

41. TR Verny: Tomorrow's baby: the art and science of parenting from conception through infancy. New York: Simon & Schuster; 2002.
42. DJ Hales, B Lozoff, R Sosa, JH Kennell: Defining the limits of the maternal sensitive period. *Dev Med Child Neurol* **19** 4 (1977), pp. 454-461.
43. P de Chateau, H Holmberg, K Jakobsson, J Winberg: A study of factors promoting and inhibiting lactation. *Dev Med Child Neurol* **19** 5 (1977), pp. 575-584.
44. P de Chateau, B Wiberg: Long-term effect on mother-infant behaviour of extra contact during the first hour post partum. III. Follow-up at one year. *Scand J Soc Med* **12** 2 (1984), pp. 91-103.
45. SK McGrath, JH Kennell: Extended mother-infant skin-to-skin contact and prospect of breastfeeding. *Acta Paediatr* **91** 12 (2002), pp. 1288-1289.
46. P de Chateau: The interaction between the infant and the environment: the importance of mother-child contact after delivery. *Acta Paediatr Scand Suppl* **344** (1988), pp. 21-30.
47. MH Klaus, R Jerauld, NC Kreger, W McAlpine, M Steffa, JH Kennel: Maternal attachment. Importance of the first post-partum days. *N Engl J Med* **286** 9 (1972), pp. 460-463.
48. P de Chateau, B Wiberg: Long-term effect on mother-infant behaviour of extra contact during the first hour post partum. II. A follow-up at three months. *Acta Paediatr Scand* **66** 2 (1977), pp. 145-151.
49. P de Chateau, B Wiberg: Long-term effect on mother-infant behaviour of extra contact during the first hour post partum. I. First observations at 36 hours. *Acta Paediatr Scand* **66** 2 (1977), pp. 137-143.
50. P de Chateau: The first hour after delivery: its impact on synchrony of the parent-infant relationship. *Paediatrician* **9** 3-4 (1980), pp. 151-168.
51. MB Hennessy: Hypothalamic-pituitary-adrenal responses to brief social separation. *Neurosci Biobehav Rev* **21** 1 (1997), pp. 11-29.
52. CL Coe, GR Lubach, ML Schneider, DJ Dierschke, WB Ershler: Early rearing conditions alter immune responses in the developing infant primate. *Pediatrics* **90** 3 Pt 2 (1992), pp. 505-509.
53. AN Schore: The effects of early relational trauma on right brain development, affect regulation, and infant mental health. *Infant Ment Health J* **22** 1-2 (2001), pp. 201-269.

54. MR Gunnar: Reactivity of the hypothalamic-pituitary-adrenocortical system to stressors in normal infants and children. *Pediatrics* **90** 3 Pt 2 (1992), pp. 491-497.
55. MR Gunnar, J Connors, J Isensee, L Wall: Adrenocortical activity and behavioral distress in human newborns. *Dev Psychobiol* **21** 4 (1988), pp. 297-310.
56. MS McClellan, WA Cabianca: Effects of early mother-infant contact following cesarean birth. *Obstet Gynecol* **56** 1 (1980), pp. 52-55.
57. G Gathwala, I Narayanan: Cesarean section and delayed contact: effect on baby's behaviour [see comments]. *Indian Pediatr* **27** 12 (1990), pp. 1295-1299.
58. G Gathwala, I Narayanan: Influence of cesarean section on mother-baby interaction. *Indian Pediatr* **28** 1 (1991), pp. 45-50.
59. MR DiMatteo, SC Morton, HS Lepper, TM Damush, MF Carney, M Pearson, KL Kahn: Cesarean childbirth and psychosocial outcomes: a meta-analysis. *Health Psychol* **15** 4 (1996), pp. 303-314.
60. P Levine: *Waking the tiger*. Berkeley: North Atlantic Books; 1997.
61. BA van der Kolk, AC McFarlane, L Weisaeth, Eds: *Traumatic stress: the effects of overwhelming experience on mind, body, and society*. New York: Guilford; 1996.
62. RM Post, SRB Weiss, M Smith: Stress, conditioning, and the temporal aspects of affective disorders. *Ann N Y Acad Sci* **771** (1995), pp. 677-695.
63. J Ledoux: *The emotional brain: the mysterious underpinnings of emotional life*. New York: Touchstone; 1996.
64. RM Post, SRB Weiss, M Smith: Sensitization and kindling: implications for the evolving neural substrate of PTSD. In: MJ Freidman, DS SCharney, AY Deutch, editors. *Neurobiology and Clinical Consequences of Stress: From Normal Adaptation to PTSD*. New York: Raven; 1995. pp. 203-244.
65. SM Antelman, AR Caggiula, S Gershon, DJ Edwards, MC Austin, S Kiss, D Kocan: Stressor induced oscillation: a possible model of the bidirectional symptoms of PTSD. *Ann N Y Acad Sci* **821** (1997), pp. 296-304.
66. SM Southwick, CA Morgan III, AD Bremner, CG Grillon, JH Krystal, LM Nagy, DS Charney: Noradrenergic alterations in Posttraumatic Stress Disorder. *Ann N Y Acad Sci* **821** (1997), pp. 125-141.
67. RM Post, DR Rubinow, JC Ballenger: Conditioning and sensitization in the longitudinal course of affective illness. *Br J Psychiatry* **149** (1986), pp. 191-201.

68. BD Perry: Memories of fear: how the brain stores and retrieves physiologic states, feelings, behaviors and thoughts from traumatic events. In: J Goodwin, R Attias, editors. *Splintered Reflections: Images of the Body in Trauma*: Basic Books; 1999.
69. RC Scaer: The whiplash syndrome: a model of traumatic stress. *The Journal of Cognitive Rehabilitation* **July/August** (2000), pp. 2-11.
70. AC McFarlane, R Yehuda: Resilience, vulnerability, and the course of posttraumatic reactions. In: BA van der Kolk, AC McFarlane, L Weisaeth, editors. *Traumatic stress: the effects of overwhelming experience on mind, body, and society*. New York: Guilford; 1996. pp. 155-181.
71. BA Campbell, J Jaynes: Reinstatement. *Psychol Rev* **73** 5 (1966), pp. 478-480.
72. BL Green: Comments on the "Empirical Basis for Biological Studies in PTSD". *Ann N Y Acad Sci* **821** (1997), pp. 52-55.
73. W Yule: Posttraumatic stress disorder in the general population and in children. *J Clin Psychiatry* **62 (Suppl 17)** (2001), pp. 23-28.
74. MM Murburg: The psychobiology of Posttraumatic Stress Disorder: an overview. *Ann N Y Acad Sci* **821** (1997), pp. 352-358.
75. BD Perry: Neurobiological sequelae of childhood trauma: Post-traumatic Stress Disorders in children. In: M Murberg, editor. *Catecholamines in Post-traumatic Stress Disorder: Emerging Concepts*. Washington, D.C.: American Psychiatric Association; 1994. pp. 253-276.
76. BD Perry: Incubated in terror: neurodevelopmental factors in the 'Cycle of Violence'. In: J Osofsky, editor. *Children, Youth and Violence: The Search for Solutions*. New York: Guilford; 1997. pp. 124-148.
77. RG Maunder, JH Hunter: Attachment and psychosomatic medicine: developmental contributions to stress and disease. *Psychosom Med* **63** (2001), pp. 556-567.
78. WT Boyce, RG Barr, LK Zeltzer: Temperament and the psychobiology of childhood stress. *Pediatrics* **90** 3 (1992), pp. 483-486.
79. BA van der Kolk, L Weisaeth, O van der Hart: History of trauma in psychiatry. In: BA van der Kolk, AC McFarlane, L Weisaeth, editors. *Traumatic stress: the effects of overwhelming experience on mind, body, and society*. New York: Guilford; 1996. pp. 47-74.

80. AY Shalev: Stress versus traumatic stress: from acute homeostatic reactions to chronic psychopathology. In: BA van der Kolk, AC McFarlane, L Weisaeth, editors. *Traumatic stress: the effects of overwhelming experience on mind, body, and society*. New York: Guilford; 1996. pp. 77-101.
81. CA Sandman, PD Wadhwa, C Dunkel-Schetter, A Chicz-DeMet, J Belman, M Porto, Y Murata, TJ Garite, FM Crinella: Psychobiological influences of stress and HPA regulation on the human fetus and infant birth outcomes. *Ann N Y Acad Sci* **739** (1994), pp. 198-210.
82. R Yehuda, EL Giller, SM Southwick, MT Lowy, JW Mason: Hypothalamic-pituitary-adrenal dysfunction in posttraumatic stress disorder. *Biol Psychiatry* **30** (1991), pp. 1030-1048.
83. CA Sandman, PD Wadhwa, A Chicz-DeMet, C Dunkel-Schetter, M Porto: Maternal stress, HPA activity, and fetal/infant outcome. *Ann N Y Acad Sci* **814** (1997), pp. 266-725.
84. P Wadhwa, CA Sandman, M Porto, C Dunkel-Schetter, TJ Garite: The association between prenatal stress and infant birth weight and gestational age at birth: a prospective investigation. *Am J Obstet Gynecol* **169** 4 (1993), pp. 858-865.
85. RL Copper, RL Goldenberg: The preterm prediction study: maternal stress is associated with spontaneous preterm birth at less than thirty-five weeks gestation. *Am J Obstet Gynecol* **175** 5 (1996), pp. 1286-1292.
86. AC McFarlane: The prevalence and longitudinal course of PTSD. *Ann N Y Acad Sci* **821** (1997), pp. 10-23.
87. A Calobrisi: Biopsychosocial study of diabetes mellitus. *Psychother Psychosom* **39** 4 (1983), pp. 193-200.
88. MB Evans: Emotional stress and diabetic control: a postulated model for the effect of emotional distress upon intermediary metabolism in the diabetic. *Biofeedback Self Regul* **10** 3 (1985), pp. 241-254.
89. P Sterling, J Eyer: Biological basis of stress-related mortality. *Soc Sci Med [E]* **15** 1 (1981), pp. 3-42.
90. G Soltesz, S Jeges, G Dahlquist: Non-genetic risk determinants for type 1 (insulin-dependent) diabetes mellitus in childhood. Hungarian Childhood Diabetes Epidemiology Study Group. *Acta Paediatr* **83** 7 (1994), pp. 730-735.
91. TF Treuting: The role of emotional factors in the etiology and course of diabetes mellitus: a review of the recent literature. *Am J Med Sci* (1962), pp. 131/193-147/109.

92. IA Mirsky: Emotional factors in the patient with diabetes mellitus. *Bull Menninger Clin* **12** (1948), pp. 187-193.
93. RS Surwit, MS Schneider: Role of stress in the etiology and treatment of diabetes mellitus. *Psychosom Med* **55** 4 (1993), pp. 380-393.
94. GE Daniels: The role of emotion in the onset and course of diabetes. *Psychosom Med* **10** (1948), pp. 288-290.
95. J Pickup, G Williams, Eds: Textbook of diabetes. 2nd ed. Osney Mead, Oxford: Blackwell Science; 1997.
96. MR Cammarn: Trauma and diabetes. *Postgrad Med* **35** (1964), pp. A62-70.
97. LEJ Hinkle, FM Evans, E Wolf, G Conger, CJ Edwards, B Pugh: IV: Life history of three persons with relatively mild, stable diabetes, and relation of significant experiences in their lives to the onset and course of the disease. *Psychosom Med* **13** 3 (1951), pp. 184-202.
98. GE Daniels: Present trends in the evaluation of psychic factors in diabetes mellitus: a critical review of experimental, general medical and psychiatric literature of the last five years. *Psychosom Med* **1** (1939), pp. 527-552.
99. EI Falstein, I Judas: Juvenile diabetes and its psychiatric implications. *Am J Orthopsychiatry* **25** (1955), pp. 330-342.
100. BR Gendel, JE Benjamin: Psychogenic factors in the etiology of diabetes. *N Engl J Med* **234** (1946), pp. 556-560.
101. G Dahlquist, L Blom, G Lonnberg: The Swedish Childhood Diabetes Study: a multivariate analysis of risk determinants for diabetes in different age groups. *Diabetologia* **34** 10 (1991), pp. 757-762.
102. WR Carter, J Herrman, K Stokes, DJ Cox: Promotion of diabetes onset by stress in the BB rat. *Diabetologia* **30** 8 (1987), pp. 674-675.
103. WK Halford, S Cuddihy, RH Mortimer: Psychological stress and blood glucose regulation in type 1 diabetic patients. *Health Psychol* **9** 5 (1990), pp. 516-528.
104. LEJ Hinkle, FM Evans, S Wolf: III: Life history of three persons with relatively labile diabetes, and relation of significant experiences in their lives to the onset and course of the disease. *Psychosom Med* **13** (1951), pp. 160-183.

105. RS Surwit, MS Schneider, MN Feinglos: Stress and diabetes mellitus. *Diabetes Care* **15** 10 (1992), pp. 1413-1422.
106. LA Gonder-Frederick, WR Carter, DJ Cox, WL Clarke: Environmental stress and blood glucose change in insulin-dependent diabetes mellitus. *Health Psychol* **9** 5 (1990), pp. 503-515.
107. A Riazi, C Bradley: Diabetes, type I. In: G Fink, editor. Encyclopedia of stress. Volume I. San Diego: Academic Press; 2000. pp. 688-693.
108. B Stabler, MA Morris, J Litton, MN Feinglos, RS Surwit: Differential glycemc response to stress in type A and type B individuals with IDDM [letter]. *Diabetes Care* **9** 5 (1986), pp. 550-552.
109. G Dahlquist: Non-genetic risk determinants of type 1 diabetes. *Diabetes Metab* **20** 3 (1994), pp. 251-257.
110. B Hagglof, L Blom, G Dahlquist, G Lonnberg, B Sahlin: The Swedish childhood diabetes study: indications of severe psychological stress as a risk factor for type 1 (insulin-dependent) diabetes mellitus in childhood. *Diabetologia* **34** 8 (1991), pp. 579-583.
111. M Rewers, RE LaPorte, M Walczak, K Dmochowski, E Bogaczynska: Apparent epidemic of insulin-dependent diabetes mellitus in Midwestern Poland. *Diabetes* **36** 1 (1987), pp. 106-113.
112. RF Hamman, EC Gay, KJ Cruickshanks, M Cook, DC Lezotte, GJ Klingensmith, HP Chase: Colorado IDDM Registry. Incidence and validation of IDDM in children aged 0-17 yr. *Diabetes Care* **13** 5 (1990), pp. 499-506.
113. G Dahlquist, L Blom, T Tuvemo, L Nystrom, A Sandstrom, S Wall: The Swedish childhood diabetes study: results from a nine year case register and a one year case-referent study indicating that type 1 (insulin-dependent) diabetes mellitus is associated with both type 2 (non-insulin-dependent) diabetes mellitus and autoimmune disorders. *Diabetologia* **32** 1 (1989), pp. 2-6.
114. PM Rothwell, SA Gutnikov, PA McKinney, E Schober, C Ionescu-Trigoviste, A Neu: Seasonality of birth in children with diabetes in Europe: multicentre cohort study. *BMJ* **319** 7214 (1999), pp. 887-888.
115. MB Oldstone, M Nerenberg, P Southern, J Price, H Lewicki: Virus infection triggers insulin-dependent diabetes mellitus in a transgenic model: role of anti-self (virus) immune response. *Cell* **65** (1991), pp. 319-331.

116. A Toniolo, T Onodera, J-W Yoon, AL Notkins: Induction of diabetes by cumulative environmental insults from viruses and chemicals. *Nature* **288** (1980), pp. 383-385.
117. MBA Oldstone: Prevention of type 1 diabetes in nonobese diabetic mice by virus infection. *Science* **239** (1988), pp. 500-502.
118. A Pundziute-Lycka, B Urbonaite, G Dahlquist: Infections and risk of type I (insulin-dependent) diabetes mellitus in Lithuanian children. *Diabetologia* **43** 10 (2000), pp. 1229-1234.
119. G Dahlquist, B Kallen: Maternal-child blood group incompatibility and other perinatal events increase the risk for early-onset type 1 (insulin-dependent) diabetes mellitus. *Diabetologia* **35** 7 (1992), pp. 671-675.
120. RJ Nelson, DL Drazen: Seasonal changes in stress responses. In: G Fink, editor. Encyclopedia of stress. Volume 3. San Diego: Academic Press; 2000. pp. 402-408.
121. SSS Lo, RYM Tun, RDG Leslie: Non-genetic factors causing type 1 diabetes. *Diabet Med* **8** (1991), pp. 609-618.
122. AN Gorsuch, KM Spencer, J Lister: Evidence for a long prediabetic period in type 1 (insulin dependent) diabetes mellitus. *Lancet* **ii** (1981), pp. 1363-1365.
123. GG Dahlquist, C Patterson, G Soltesz: Perinatal risk factors for childhood type 1 diabetes in Europe. The EURODIAB Substudy 2 Study Group. *Diabetes Care* **22** 10 (1999), pp. 1698-1702.
124. CC Patterson, DJ Carson, DR Hadden, NR Waugh, SK Cole: A case-control investigation of perinatal risk factors for childhood IDDM in northern Ireland and Scotland. *Diabetes Care* **17** 5 (1994), pp. 376-381.
125. DJ Betteridge, Ed: Diabetes: current perspectives. London: Martin Dunitz; 2000.
126. GG Dahlquist, S Ivarsson, B Lindberg, M Forsgren: Maternal enteroviral infection during pregnancy as a risk factor for childhood IDDM. A population-based case-control study. *Diabetes* **44** 4 (1995), pp. 408-413.
127. PA McKinney, R Parslow, K Gurney, G Law, HJ Bodansky, DRR Williams: Antenatal risk factors for childhood diabetes mellitus: a case control study of medical record data in Yorkshire, UK. *Diabetologia* **40** (1997), pp. 933-939.
128. G Dahlquist, G Frisk, SA Ivarsson, L Svanberg, M Forsgren, H Diderholm: Indications that maternal coxsackie B virus infection during pregnancy is a risk factor for childhood-onset IDDM. *Diabetologia* **38** 11 (1995), pp. 1371-1373.

129. G Dahlquist, SS Bennich, B Kallen: Intrauterine growth pattern and risk of childhood onset insulin dependent (type I) diabetes: population based case-control study. *BMJ* **313** 7066 (1996), pp. 1174-1177.
130. SM Heidrich, MS Cranley: Effect of fetal movement, ultrasound scans, and amniocentesis on maternal-fetal attachment. *Nurs Res* **38** 2 (1989), pp. 81-84.
131. G Dahlquist, B Kallen: Early neonatal events and the disease incidence in nonobese diabetic mice. *Pediatr Res* **42** 4 (1997), pp. 489-491.
132. HP Schobel, T Fischer, K Heuszer, H Geiger, RE Schmieder: Preeclampsia: a state of sympathetic overactivity. *N Engl J Med* **335** 20 (1996), pp. 1480-1485.
133. FP Zuspan: Catecholamines. Their role in pregnancy and the development of pregnancy-induced hypertension. *J Reprod Med* **23** 3 (1979), pp. 143-150.
134. T Abboud, R Artal, F Sarkis, EH Henriksen, RK Kammula: Sympathoadrenal activity, maternal, fetal, and neonatal responses after epidural anesthesia in the preeclamptic patient. *Am J Obstet Gynecol* **144** 8 (1982), pp. 915-918.
135. JP Greenwood, EM Scott, JJ Walker, JB Stoker, DASG Mary: The magnitude of sympathetic hyperactivity in pregnancy-induced hypertension and preeclampsia. *Am J Hypertens* **16** (2003), pp. 194-199.
136. RM Lewinsky, S Riskin-Mashiah: Autonomic imbalance in preeclampsia: evidence for increased sympathetic tone in response to the supine-pressor test. *Obstet Gynecol* **91** 6 (1998), pp. 935-939.
137. EL Ryding, B Wijma, K Wijma: Posttraumatic stress reactions after emergency cesarean section. *Acta Obstet Gynecol Scand* **76** (1997), pp. 856-861.
138. EJ Mayer, RF Hamman, EC Gay, DC Lezotte, DA Savitz, GJ Klingensmith: Reduced risk of IDDM among breast-fed children. The Colorado IDDM Registry. *Diabetes* **37** 12 (1988), pp. 1625-1632.
139. L Blom, G Dahlquist, L Nystrom, A Sandstrom, S Wall: The Swedish childhood diabetes study: social and perinatal determinants for diabetes in childhood. *Diabetologia* **32** 1 (1989), pp. 7-13.
140. JN Kostraba, KJ Cruickshanks, J Lawler-Heavner, LF Jobim, MJ Rewers, EC Gay, HP Chase, G Klingensmith, RF Hamman: Early exposure to cow's milk and solid foods in infancy, genetic predisposition, and risk of IDDM. *Diabetes* **42** 2 (1993), pp. 288-295.

141. J Karjalainen, JM martin, M Knip, J Ilonen, BH Robinson, E Savilahti, HK Akerblom, HM Dosch: A bovine serum albumin peptide as a possible trigger of insulin dependent diabetes mellitus. *N Engl J Med* **327** (1992), pp. 302-307.
142. JM Norris, B Beaty, G Klingensmith, L Yu, M Hoffman, HP Chase, HA Erlich, RF Hamman, GS Eisenbarth, M Rewers: Lack of association between early exposure to cow's milk protein and beta-cell autoimmunity. Diabetes Autoimmunity Study in the Young (DAISY) [see comments]. *JAMA* **276** 8 (1996), pp. 609-614.
143. DA Schatz, NK Maclaren: Cow's milk and insulin-dependent diabetes mellitus: innocent until proven guilty. *JAMA* **276** 8 (1996), pp. 647-648.
144. U Samuelsson, C Johansson, J Ludvigsson: Breastfeeding seems to play a marginal role in the prevention of insulin-dependent diabetes mellitus. *Diabetes Res Clin Pract* **19** (1993), pp. 203-210.
145. K Borch-Johnsen, G Joner, T Mandrup-Poulsen, M Christy, B Zachau-Christiansen, K Kastrup, J Nerup: Relation between breast-feeding and incidence of insulin-dependent diabetes mellitus: a hypothesis. *Lancet* **2** (1984), pp. 1083-1086.
146. Z Huang, V Cabanela, T Howell: Stress, bottlefeeding, and diabetes [letter; comment]. *Lancet* **350** 9081 (1997), pp. 889.
147. M Karvonen, M Viik-Kajander, E Moltchanova, I Libman, R LaPorte, J Tuomilehto: Incidence of childhood type 1 diabetes worldwide. Diabetes Mondiale (DiaMond) Project Group. *Diabetes Care* **23** 10 (2000), pp. 1516-1526.
148. WJ Riley, NK Maclaren, J Krischer, R Spillar, J Silverstein, DA Schatz, S Schwartz, J Malone, S Shah, C Vadheim and others: A prospective study of the development of diabetes in relatives of patients with insulin-dependent diabetes. *N Engl J Med* **323** 17 (1990), pp. 1167-1172.
149. J Lawler-Heavner, KJ Cruickshanks, WW Hay, EC Gay, RF Hamman: Birth size and risk of insulin-dependent diabetes mellitus (IDDM). *Diabetes Res Clin Pract* **24** 3 (1994), pp. 153-159.
150. C Johnston, BA Milward, P Hoskins, RDG Leslie, GF Bottazzo, DA Pyke: Islet-cell antibodies as predictors of the later development of type 1 (insulin-dependent) diabetes. A study in identical twins. *Diabetologia* **32** (1989), pp. 382-386.
151. E Bonifacio, PJ Bingley, M Shattock, BM Dean, D Dunger, EAM Gale, GF Bottazzo: Quantification of islet-cell antibodies and prediction of insulin-dependent diabetes. *Lancet* **335** (1990), pp. 147-149.
152. DR Wegmann, GS Eisenbarth: It's insulin. *J Autoimmun* **15** (2000), pp. 286-291.

153. DG Leslie, RB Elliot: Early environmental events as a cause of IDDM. *Diabetes* **43** (1994), pp. 843-850.
154. M Vives-Pi, N Somoza, F Vargas, P Armengol, Y Sarri, JY Wu, R Pujol-Borrell: Expression of glutamic acid decarboxylase (GAD) in the alpha, beta and delta cells of normal and diabetic pancreas: implications for the pathogenesis of type I diabetes. *Clin Exp Immunol* **92** 3 (1993), pp. 391-396.
155. L Yu, M Rewers, R Gianani, E Kawasaki, Y Zhang, C Verge, P Chase, G Klingensmith, H Erlich, J Norris and others: Antiislet autoantibodies usually develop sequentially rather than simultaneously. *J Clin Endocrinol Metab* **81** 12 (1996), pp. 4264-4267.
156. J Yu, L Yu, TL Bugawan, HA Erlich, K Barriga, M Hoffman, M Rewers, GS Eisenbarth: Transient antiislet autoantibodies: infrequent occurrence and lack of association with "genetic" risk factors. *J Clin Endocrinol Metab* **85** 7 (2000), pp. 2421-2328.
157. KM Spencer, A Tarn, BM Dean, J Lister, GF Bottazzo: Fluctuating islet-cell autoimmunity in unaffected relatives of patients with insulin-dependent diabetes. *Lancet* **1** (1984), pp. 764-766.
158. L Yu, DT Robles, N Abiru, P Kaur, M Rewers, K Kelemen, GS Eisenbarth: Early expression of antiinsulin autoantibodies of humans and the NOD mouse: evidence for early determination of subsequent diabetes. *Proc Natl Acad Sci U S A* **97** 4 (2000), pp. 1701-1706.
159. RM Sapolsky: Why zebras don't get ulcers: an updated guide to stress, stress-related diseases, and coping. New York: W.H. Freeman & Co; 1998.
160. JK Karjalainen: Islet cell antibodies as predictive markers for IDDM in children with high background incidence of disease. *Diabetes* **39** (1990), pp. 1146-1150.
161. C Greenbaum, K Sears, S Kahn, J Palmer: Relationship of B-cell function and autoantibodies to progression and nonprogression of subclinical type I diabetes. *Diabetes* **48** (1999), pp. 170-175.
162. G Goddard, D McIntyre, C Leetch: A permanent change in brain functioning resulting from daily electrical stimulation. *Exp Neurol* **25** (1969), pp. 295-330.